


Lesson 10

A-Level Pure Mathematics : Year 2 Differentiation III

10.1 Differentiating $x = f(y)$

The graph is of the curve with equation $x = e^y \cos y$ with y in radians.



Video : <http://www.NumberWonder.co.uk/v9028/10a.mp4>


- (i) Obtain an expression for $\frac{dy}{dx}$ in terms of y

- (ii) What is the equation of the normal to the curve when $y = 0$?

- (iii) Add your part (ii) normal onto the graph above

[2, 2, 2 marks]

10.2 Differentiating $\arcsin x$



The graph of $y = \arcsin x$

It is the inverse of a one-to-one piece of the $\sin x$ function

$$y = \arcsin x \Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$$

Proof

Teaching Video : <http://www.NumberWonder.co.uk/v9028/10b.mp4>



Watch the video and
then write out the
proof here




[6 marks]

10.3 Exercise

Marks Available : 40

Question 1

The graph is of the curve with equation $y = \frac{\sin(x^2)}{x}$ with x in radians.



- (i) Obtain an expression for $\frac{dy}{dx}$ in terms of x

[3 marks]


- (ii) What is the *exact* equation of the tangent to the curve when $x = \sqrt{\pi}$

[5 marks]

- (iii) Add your part (ii) tangent onto the graph above

[1 mark]

Question 2



The graph of $y = \arccos x$

It is the inverse of a one-to-one piece of the $\cos x$ function


$$y = \arccos x \Rightarrow \frac{dy}{dx} = -\frac{1}{\sqrt{1-x^2}}$$

Assuming standard results for $\sin x$ and $\cos x$ prove the above result.

[7 marks]

Question 3

The graph is of a piece of the curve with equation $y = x^2 \sin x$



- (i) Obtain an expression for $\frac{dy}{dx}$ in terms of x

[3 marks]


- (ii) What is the *exact* equation of the normal to the curve when $x = \pi$

[6 marks]

- (iii) Add your part (ii) normal onto the graph above

[1 mark]

Question 4



The graph of $y = \arctan x$

It is the inverse of a one-to-one piece of the $\tan x$ function

$$y = \arctan x \Rightarrow \frac{dy}{dx} = \frac{1}{1 + x^2}$$

- (i) Show that if $y = \tan x$ then $\frac{dy}{dx} = \sec^2 x$ by using the derivatives of $\sin x$ and $\cos x$ and the quotient rule.


[2 marks]

- (ii) Hence prove that if $y = \arctan x$ then $\frac{dy}{dx} = \frac{1}{1 + x^2}$

[6 marks]

Question 5

A curve has equation, $y = \cos^2 x + \sin x$ $0 < x < 2\pi$
Find the coordinates of its stationary points.



[6 marks]

This document is a part of a **Mathematics Community Outreach Project** initiated by Shrewsbury School

It may be freely duplicated and distributed, unaltered, for non-profit educational use

In October 2020, Shrewsbury School was voted "**Independent School of the Year 2020**"

© 2022 Number Wonder

Teachers may obtain detailed worked solutions to the exercises by email from mhh@shrewsbury.org.uk