

Lesson 3

A-Level Applied Mathematics Mechanics : Moments : Year 2


3.1 On the Edge Of Tipping

Example

A 250 kg mass is bolted onto the end of a uniform plank of wood AB of length 8 m and mass 45 kg.

The plank rests on two supports at A and C where $AC = 3$ m.

The plank is in equilibrium and horizontal.



Teaching Video : [http://www.NumberWonder.co.uk/Video/v9069\(3\).mp4](http://www.NumberWonder.co.uk/Video/v9069(3).mp4)

- (a) By taking moments about A determine the reaction force at C .

- (b) By equating the total force upwards with the total force downward determine the reaction force at A .


The plank of wood is being used as a diving board.

- (c) By taking moments about C , determine the maximum mass that can be at B with the plank remaining in equilibrium ?

- (d) In the light of your previous answers, what advice would you give regarding the safe use of this diving board ?

3.2 Exercise

Question 1



A uniform plank of mass 100 kg and length 10 m rests horizontally on two smooth supports, A and B , as shown. A man of mass 80 kg starts walking from one end of the plank, A , to the other end. When he is a distance x past B the plank starts to tip.

- (i) Draw a diagram showing all significant forces on the plank when it is on the verge of tipping.

[2 marks]

- (ii) When the plank is about to tip, what will be the reaction at A ?

[1 mark]

- (iii) When the plank is about to tip, what will be the reaction at B ?


[1 mark]

- (iv) Find the distance, x , the man can walk past B before the plank starts to tip.

[4 marks]

Question 2


A uniform rod of mass 14 kg rests on two supports as shown below.



- (a) The reaction at the left hand support is 75 newtons.
What, in newtons, is the reaction at the other support ?
- (b) A mass, m , is now placed an the right hand end of the rod, such that the rod is on the point of tipping about the right hand support.
- (i) What, now, is the reaction at the left hand support ?
- (ii) What, in kg, is the mass, m ?
- (iii) What, in N, is the reaction at the right hand support ?

Question 3

M1 Exam question, 12th January 2007, Q2



A uniform plank AB has weight 120 N and length 3 m .

The plank rests horizontally in equilibrium on two smooth supports C and D , where $AC = 1\text{ m}$ and $CD = x\text{ m}$.

The reaction of the support on the plank at D has magnitude 80 N .

Modelling the plank as a rod,

(a) show that $x = 0.75$

[3 marks]

A rock is now placed at B and the plank is on the point of tilting about D

Modelling the rock as a particle, find

(b) the weight of the rock

[4 marks]

(c) the magnitude of the reaction of the support on the plank at D


[2 marks]

(d) State how you have used the model of the rock as a particle

[1 mark]

Question 4

A-Level Examination Question from January 2019, IAL, M1, Q4



A boy sees a box on the end Q of a plank PQ which overhangs a swimming pool. The plank has mass 30 kg, is 5 m long and rests in a horizontal position on two bricks. The bricks are modelled as smooth supports, one acting on the plank at P and one acting on the plank at R , where PR is 3 m. The support at R is on the edge of the swimming pool, as shown. The boy has a mass 40 kg and the box has mass 2.5 kg. The plank is modelled as a uniform rod and the boy and the box as particles.

The boy steps on to the plank at P and begins to walk slowly along the plank towards the box.

- (a) Find the distance he can walk along the plank from P before the plank starts to tilt.

[4 marks]

- (b) State how you have used, in your working, the fact that the box is modelled as a particle.

[1 mark]

A rock of mass M kg is placed on the plank at P . The boy is then able to walk slowly along the plank to the box at the end Q without the plank tilting. The rock is modelled as a particle.

- (c) Find the smallest possible value of M .

[4 marks]