

## Lesson 8

### A-Level Pure Mathematics, Year 2 Functions II

#### 8.1 The Möbius Function

A Möbius function is of the form,

$$f(x) = \frac{ax + b}{cx + d} \quad x \in \mathbb{R}, \quad ad - bc \neq 0, \quad x \neq -\frac{d}{c}$$

where  $a$ ,  $b$ ,  $c$  and  $d$  are constants.

Finding the inverse of this one-to-one function requires a method for dealing with the fact that the  $x$  appears in two different places, in both numerator and denominator.

#### Example

(i) Find the inverse of,

$$h(x) = \frac{5x + 2}{3x + 4} \quad x \in \mathbb{R}, \quad x \neq -\frac{4}{3}$$

[ 4 marks ]

(ii) State the domain of the inverse function.

[ 1 mark ]

## 8.2 Exercise

*Any solution based entirely on graphical  
or numerical methods is not acceptable*

Marks Available: 54

### Question 1

- ( i ) Find the inverse of,  $m(x) = \frac{7x + 1}{2x + 3}$   $x \in \mathbb{R}, x \neq -\frac{3}{2}$

[ 4 marks ]

- ( ii ) State the domain of the inverse function

[ 1 mark ]

**Question 2**

Prove that the inverse of

$$f(x) = \frac{ax + b}{cx + d} \quad x \in \mathbb{R}, \quad ad - bc \neq 0, \quad x \neq -\frac{d}{c}$$

where  $a, b, c$  and  $d$  are constants, is

$$f^{-1}(x) = \frac{-dx + b}{cx - a} \quad x \in \mathbb{R}, \quad x \neq \frac{a}{c}$$

[ 6 marks ]

### Question 3


- ( i ) Find the inverse of,  $s(x) = \frac{2x - 1}{x - 2} \quad x \neq 2$   
by using the result proven in question 2

[ 2 marks ]

- ( ii ) This is an example of a “self inverse” function.  
Explain what this means.

[ 1 mark ]

- ( iii ) Plot an accurate graph of  $s(x)$  and also the line  $y = x$   
Mark on the horizontal and the vertical asymptotes of  $s(x)$



[ 3 marks ]

- ( iv ) What property do the graphs of all self-inverse functions have ?

[ 2 marks ]

**Question 4**

- ( i ) Write down the equations of the two asymptotes of the function,

$$f(x) = \frac{2}{x} \quad x \in \mathbb{R}, \quad x \neq 0$$

[ 2 marks ]

- ( ii ) With your part (i) answer in mind, write down the equations of the two asymptotes of the graph of the function,

$$h(x) = \frac{2}{x+1} + 3$$

[ 2 marks ]

- ( iii ) What number must be excluded from the domain of  $h(x)$  ?

[ 1 mark ]

- ( iv ) Sketch the graph of  $h(x)$   
(No need for an accurate graph, just the essential shape and the asymptotes)

[ 3 marks ]

( v ) Show that  $h(x)$  is a Möbius transformation by writing it in the form,

$$f(x) = \frac{ax + b}{cx + d} \quad x \in \mathbb{R}, \quad ad - bc \neq 0, \quad x \neq -\frac{d}{c}$$

where  $a, b, c$ , and  $d$  are constants the values of which you have determined.


[ 3 marks ]

( vi ) Find  $h^{-1}(x)$ , stating its domain.

[ 4 marks ]

**Question 5**

*A-Level Examination Question from November 2018, Paper C34, Q10 (Edexcel)*



The sketch is of the graph with equation  $y = g(x)$ , where,

$$g(x) = \frac{3x - 4}{x - 3}, \quad x \in \mathbb{R}, \quad x < 3$$

The graph cuts the  $x$ -axis at the point  $A$  and the  $y$ -axis at the point  $B$ , as shown.

- (a) State the range of  $g$  [ 1 mark ]
- (b) State the coordinates of,
- (i) point  $A$  [ 1 mark ]
- (ii) point  $B$  [ 1 mark ]
- (c) Find  $gg(x)$  in its simplest form.

[ 3 marks ]

- ( d ) Sketch the graph with equation  $y = |g(x)|$   
On your sketch, show the coordinates of each point at which the graph meets or cuts the axes and state the equation of each asymptote.

[ 3 marks ]

- ( e ) Find the exact solution of the equation  $|g(x)| = 8$


[ 3 marks ]



### Question 6

$$f(x) = |x - 2| + 1, \quad x \in \mathbb{R}$$

The function  $f(x)$  has a “critical value” when  $|x - 2| = 0$  in the sense that there will be a “drastic change” in the smoothness of the graph at  $x = 2$



This function can also be written in hybrid form as,  $f(x) = \begin{cases} x - 1 & x \geq 2 \\ -x + 3 & x < 2 \end{cases}$

Consider now the following function,

$$g(x) = |x + 1| + |2x + 1| - |x - 2|$$

Determine the critical values of  $g(x)$ , sketch  $g(x)$  and express  $g(x)$  in hybrid form.

[ 8 marks ]