

3.1 Cubic and Roots

Having taken a fresh look at quadratic equations through a study of their roots, the focus can now shift to cubic equations. Thanks to Gerolamo Cardano it is known that there is, in principle, a formula to solve cubics. Alas, it has a square root nested inside a cube root which makes manipulating it intricate. Fortunately, the algebra can be skirted around by a more thoughtful approach. To illustrate the idea, it will first be applied to the generalised quadratic equation.

3.2 A Proof Rewritten

The Roots of a Quadratic

If α and β are the roots of the equation $ax^2 + bx + c = 0$ $a, b, c \in \mathbb{C}$

then, $\alpha + \beta = -\frac{b}{a}$ $\alpha\beta = \frac{c}{a}$

Proof

For a general quadratic, with roots α and β ,

$$\begin{aligned} ax^2 + bx + c &= a\left(x^2 + \frac{b}{a}x + \frac{c}{a}\right) \\ &= a(x - \alpha)(x - \beta) \\ &= a(x^2 - \alpha x - \beta x + \alpha\beta) \\ &= a(x^2 - (\alpha + \beta)x + \alpha\beta) \end{aligned}$$

From which can be seen that,

$$\left(x^2 + \frac{b}{a}x + \frac{c}{a}\right) = (x^2 - (\alpha + \beta)x + \alpha\beta)$$

And, by matching coefficients of x , the deduction made that,

$$\alpha + \beta = -\frac{b}{a} \quad \text{and} \quad \alpha\beta = \frac{c}{a} \quad \square$$

The key feature of this proof, in comparison with that presented in Lesson 1, is that no use is made of the formula,

$$x = \frac{-b \pm \sqrt{b^2 + 4ac}}{2a}$$

3.3 A Similar Proof For Cubics

The Roots of a Cubic

If α, β and γ are roots of $ax^3 + bx^2 + cx + d = 0$ $a, b, c, d \in \mathbb{C}$

$$\text{then, } \alpha + \beta + \gamma = -\frac{b}{a} \quad \alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a} \quad \alpha\beta\gamma = -\frac{d}{a}$$

Proof

For a general cubic, with roots α, β and γ ,

$$\begin{aligned} ax^3 + bx^2 + cx + d &= a \left(x^3 + \frac{b}{a}x^2 + \frac{c}{a}x + \frac{d}{a} \right) \\ &= a \left((x - \alpha)(x - \beta)(x - \gamma) \right) \\ &= a \left((x^2 - \alpha x - \beta x + \alpha\beta)(x - \gamma) \right) \\ &= a \left((x^2 - \alpha x - \beta x + \alpha\beta)x - (x^2 - \alpha x - \beta x + \alpha\beta)\gamma \right) \\ &= a \left(x^3 - \alpha x^2 - \beta x^2 + \alpha\beta x - \gamma x^2 + \alpha\gamma x + \beta\gamma x - \alpha\beta\gamma \right) \\ &= a \left(x^3 - (\alpha + \beta + \gamma)x^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)x - \alpha\beta\gamma \right) \end{aligned}$$

from which can be seen that,


$$\left(x^3 + \frac{b}{a}x^2 + \frac{c}{a}x + \frac{d}{a} \right) = \left(x^3 - (\alpha + \beta + \gamma)x^2 + (\alpha\beta + \beta\gamma + \gamma\alpha)x - \alpha\beta\gamma \right)$$

and, by matching coefficients of x , the deduction made that,

$$\alpha + \beta + \gamma = -\frac{b}{a}, \quad \alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a} \quad \text{and} \quad \alpha\beta\gamma = -\frac{d}{a} \quad \square$$

Notice that, as with the quadratic,

- The sum of the roots is $\left(-\frac{b}{a}\right)$
- The sum of all possible product pairs of roots is $\left(\frac{c}{a}\right)$



3.4 Example

The roots of the cubic equation $2x^3 + 5x^2 - 2x + 3 = 0$ are α , β and γ .

Without solving the equation, find the value of $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$

Teaching Video : <http://www.NumberWonder.co.uk/v9093/3.mp4>



Watch the teaching video and then write out a solution to the question.



[4 marks]

3.5 Exercise

*Any solution based entirely on graphical
or numerical methods is not acceptable.*

Make the method used clear.

Marks available : 40

Question 1

The roots of the equation $4x^3 - 3x^2 - x + 6 = 0$ are α, β and γ

Without solving the equation, find the roots of

(i) $\alpha + \beta + \gamma$

(ii) $\alpha\beta + \beta\gamma + \gamma\alpha$

[1, 1 mark]

(iii) $\alpha^2 \beta^2 \gamma^2$

(iv) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$

[1, 2 marks]

Question 2

The roots of the cubic equation $ax^3 + bx^2 + cx + d = 0$ are,

$$\alpha = 8, \quad \beta = 9 \quad \text{and} \quad \gamma = -10$$

Find integer values for a, b, c and d

[2 marks]

Question 3

	α	β	γ	
α	α^2	$\alpha\beta$	$\alpha\gamma$	$\alpha + \beta + \gamma$
β	$\alpha\beta$	β^2	$\beta\gamma$	
γ	$\alpha\gamma$	$\beta\gamma$	γ^2	
	$\alpha + \beta + \gamma$			

- (i) With the aid of the diagram expand the brackets of $(\alpha + \beta + \gamma)^2$

[1 mark]

The roots of the equation $2x^3 + 4x^2 + 7x + 1 = 0$ are α, β and γ

Without solving the equation, write down the values of,

- (ii) $\alpha + \beta + \gamma$

[1 mark]

- (iii) $\alpha\beta + \beta\gamma + \gamma\alpha$

[1 mark]

- (iv) $\alpha\beta\gamma$

[1 mark]

- (v) $\alpha^2 + \beta^2 + \gamma^2$

[2 marks]

Question 4

Further A-Level Examination Question from June 2017 SAM, Core 2, Q1

The roots of the equation $x^3 - 8x^2 + 28x - 32 = 0$ are α, β and γ

Without solving the equation, find the value of

(i) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$

[3 marks]

(ii) $(\alpha + 2)(\beta + 2)(\gamma + 2)$

[3 marks]

(iii) $\alpha^2 + \beta^2 + \gamma^2$

[2 marks]

Question 5

Our sequence of questions exploring Cardano's extraordinary achievement in finding and using a formula to solve cubic equations continues by looking at,

$$x^3 - 252x + 1296 = 0$$

As you will discover, this example involves complex numbers.

Here is a recap of the theory to be applied;

Roots of a Cubic

Given a depressed cubic of the form

$$t^3 + pt + q = 0 \quad p, q \in \mathbb{C}$$

where p and q are not both zero, and $4p^3 + 27q^2 \neq 0$, calculate,

$$C = \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}$$

A root of the cubic is then given by,

$$\alpha = C - \frac{p}{3C}$$

Polynomial division can then be used to find the remaining roots.

To help with a tricky step, there is a part (i) which is useful when tackling the main problem.

(i) Expand the brackets,

$$(6 + 4\sqrt{3}i)^3 \text{ where } i = \sqrt{-1} \text{ such that } i^2 = -1$$

giving your answer in the form $a\sqrt{3}i + b$ for integer a and b

[3 marks]

- (ii) Determine using the method of Cardano, the real root of the cubic,

$$x^3 - 252x + 1296 = 0$$


[3 marks]

- (iii) Using polynomial division with your part (ii) answer, find all three roots, which are all real, of the depressed cubic,

$$x^3 - 252x + 1296 = 0$$

[2 marks]

Question 6



- (i) In question 3 a square with sides of length $\alpha + \beta + \gamma$ was of assistance in expanding the brackets of $(\alpha + \beta + \gamma)^2$

For $(\alpha + \beta + \gamma)^3$ a cube can be used in which case the three layers of the cube give cuboids with volumes as shown above.

By using the diagrams or otherwise, expand the brackets of $(\alpha + \beta + \gamma)^3$

[1 mark]

- (ii) Manipulate your part (i) answer into a form that will allow you to express $\alpha^3 + \beta^3 + \gamma^3$ in terms of $\alpha + \beta + \gamma$, $\alpha\beta + \beta\gamma + \gamma\alpha$ and $\alpha\beta\gamma$

[2 marks]

Question 7

Further A-Level Examination Question from June 2019, Core 2, Q2 (Edexcel)

The roots of the equation $x^3 - 2x^2 + 4x - 5 = 0$ are p, q and r

Without solving the equation, find the value of

(i) $\frac{2}{p} + \frac{2}{q} + \frac{2}{r}$

[3 marks]

(ii) $(p - 4)(q - 4)(r - 4)$

[3 marks]

(iii) $p^3 + q^3 + r^3$

[2 marks]

This document is a part of a **Mathematics Community Outreach Project** initiated by Shrewsbury School

It may be freely duplicated and distributed, unaltered, for non-profit educational use

In October 2020, Shrewsbury School was voted “**Independent School of the Year 2020**”

© 2025 Number Wonder

Teachers may obtain detailed worked solutions to the exercises by email from MHHShrewsbury@Gmail.com