Further Pure A-Level Mathematics Compulsory Course Component Core 2

TELESCOPING SERIES



THE METHOD OF DIFFERENCES

TELESCOPING SERIES

THE METHOD OF DIFFERENCES

Lesson 1

Further A-Level Pure Mathematics, Core 2

Telescoping Series

1.1 The Classic Telescoping Series Example

Often in mathematics, a seemingly unremarkable question can lead to an ingenious solution with an underlying technique, then of use in solving many similar and seeming harder questions. This in turn leads to a focus on questions that one feels should be solvable in a likewise manner but which are not.

A excellent example of such an apparently unremarkable question is the sum,

$$\sum_{r=1}^{n} \frac{1}{r(r+1)} = \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{n(n+1)}$$

One approach is to look at the initial partial sums of the series,

$$\frac{1}{2} = \frac{1}{2}$$

$$\frac{1}{2} + \frac{1}{6} = \frac{2}{3}$$

$$\frac{1}{2} + \frac{1}{6} + \frac{1}{12} = \frac{3}{4}$$

$$\frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} = \frac{4}{5}$$

$$\frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \frac{1}{30} = \frac{5}{6}$$

A reasonable guess at this stage would be that,


$$\sum_{r=1}^{n} \frac{1}{r(r+1)} = \frac{n}{n+1}$$

This guess could be proven correct with a proof by induction.

There is an unsatisfactory messiness with this approach for it does not give insightful about why the result is true.

For the "insight" an old tool is needed, that of partial fractions, along with a willingness to explore the resulting equivalent series.

Teaching Video: http://www.NumberWonder.co.uk/v9097/1a.mp4
http://www.NumberWonder.co.uk/v9097/1b.mp4



After watching the video, write out the details of the Telescoping Series method of showing that,

$$\sum_{r=1}^{n} \frac{1}{r(r+1)} = \frac{n}{n+1}$$

1.2 Exercise

Any solution based entirely on graphical or numerical methods is not acceptable

Marks Available: 34

Question 1

(a) Express
$$\frac{1}{(r+2)(r+3)}$$
 in partial fractions

[1 mark]

(**b**) Hence use the Telescoping Series method to find the sum of the series,

$$\sum_{r=1}^{n} \frac{1}{(r+2)(r+3)}$$

The Telescoping Series are the result of applying a technique known as

The Method of Differences

Question 2

Further A-Level Examination Question from June 2010, FP2, Q1

(a) Express
$$\frac{3}{(3r-1)(3r+2)}$$
 in partial fractions

[2 marks]

(**b**) Using your answer to part (a) and the method of differences, show that

$$\sum_{r=1}^{n} \frac{3}{(3r-1)(3r+2)} = \frac{3n}{2(3n+2)}$$

[3 marks]

(c) Evaluate
$$\sum_{r=100}^{1000} \frac{3}{(3r-1)(3r+2)}$$
 giving your answer to 3 significant figures

Question 3

(a) Express $\frac{1}{r(r+2)}$ in partial fractions

[1 mark]

(**b**) Hence find the sum of the series $\sum_{r=1}^{n} \frac{1}{r(r+2)}$ using the method of differences

Question 4

Further A-Level Examination Question from June 2018, IAL, F2, Q5

(a) Express
$$\frac{4r+2}{r(r+1)(r+2)}$$
 in partial fractions

[3 marks]

(**b**) Hence using the method of differences, prove that

$$\sum_{r=1}^{n} \frac{4r+2}{r(r+1)(r+2)} = \frac{n(an+b)}{2(n+1)(n+2)}$$

where a and b are constants to be found

Question 5

Further A-Level Examination Question from June 2017, FP2, Q1

(a) Show that, for r > 0

$$\frac{1}{r^2} - \frac{1}{(r+1)^2} \equiv \frac{2r+1}{r^2(r+1)^2}$$

[1 mark]

(**b**) Hence prove that, for $n \in \mathbb{N}$

$$\sum_{r=1}^{n} \frac{2r+1}{r^{2}(r+1)^{2}} = \frac{n(n+2)}{(n+1)^{2}}$$

(c) Show that, for $n \in \mathbb{N}$, n > 1

$$\sum_{r=n}^{3n} \frac{6r+3}{r^2(r+1)^2} = \frac{an^2+bn+c}{n^2(3n+1)^2}$$

where a, b and c are constants to be found.

[3 marks]