

Lesson 10

Further A-Level Pure Mathematics, Core 2 Hyperbolic Functions

10.1 By Parts and Substitution

This lesson is a revision of the two topics Integration by Parts and Integration by Substitution but with an emphasis on situations that involve the hyperbolic functions and their associated trigonometric identities.

Keep in mind that carrying out an integration using parts (LIDI) or a substitution can be a lot of work and there is a definite time saving in opting for a “chain rule backwards” if a question is amenable to that approach. However, an examination question may prescribe that the solution method has to be “by parts” or “by substitution”, or the parts or substitution methods may genuinely be the easier method of doing the question.

10.2 Example #1 (By Parts)

Use integration by parts to find $\int x \sinh 3x \, dx$

[3 marks]

10.3 Example #2 (Trigonometric Odd Powers)

In general to integrate a trigonometric function raised to an odd power the following simple manoeuvre can be very effective.

Trigonometric Odd Powers Integration

For small odd values of n , use the cunning moves,

$$\int \sinh^n x \, dx = \int \sinh x \sinh^{n-1} x \, dx$$

$$\int \cosh^n x \, dx = \int \cosh x \cosh^{n-1} x \, dx$$

Find $\int \sinh^3 x \, dx$

[3 marks]

10.4 Example #3 (Substitution)

When carrying out an integration involving a substitution $x = f(u)$ remember that three items need attending to:

- Making the substitution into the function being integrated
- Replacing the dx with du having first worked out $\frac{dx}{du}$
- Replacing the “ $x =$ ” limits with “ $u =$ ” limits

Use the substitution $x = 3 \sinh u$ to find the exact value of $\int_0^6 \frac{x^3}{\sqrt{x^2 + 9}} dx$

10.5 Exercise

*Any solution based entirely on graphical
or numerical methods is not acceptable*
Marks Available : 30

Question 1

The graph is of the function $f(x) = \cos^3 x$, $0 \leq x \leq \pi$



Find the area shaded which is bounded by $f(x)$ and the positive coordinate axes.

[5 marks]

Question 2

Use integration by parts to find $\int 25x^2 \cosh(5x) dx$

[5 marks]

Question 3

Find $\int \cosh^3 x dx$

[3 marks]

Question 4

Use the substitution $x = 6 \cosh u$ to find the exact value of $\int_{12}^{18} \frac{x^3}{\sqrt{x^2 - 36}} dx$

Give your answer in the form $s(p\sqrt{2} - q\sqrt{3})$ where s is a square number and p and q are prime numbers.

[11 marks]

Question 5

Using the substitution $x = \sec \theta$ find,

(i) $\int \frac{1}{x \sqrt{x^2 - 1}} dx$

[2 marks]

(ii) $\int \frac{\sqrt{x^2 - 1}}{x} dx$

[4 marks]

This document is a part of a **Mathematics Community Outreach Project** initiated by Shrewsbury School

It may be freely duplicated and distributed, unaltered, for non-profit educational use

In October 2020, Shrewsbury School was voted “**Independent School of the Year 2020**”

© 2025 Number Wonder

Teachers may obtain detailed worked solutions to the exercises by email from MHHShrewsbury@Gmail.com