

Lesson 8

Further A-Level Pure Mathematics, Core 2 Hyperbolic Functions

8.1 Integration (Chain Rule Backwards)

In principle, given we've already[†] looked at differentiation questions, tackling an exercise on integration is just a case of using the following, previously supplied, table backwards, and remembering that if it's an indefinite integral (one without limits) to put in the constant of integration, c .

$f(x)$	$f'(x)$	In Formula Book ?
$\sinh x$	$\cosh x$	Yes
$\cosh x$	$\sinh x$	Yes
$\tanh x$	$\operatorname{sech}^2 x$	Yes
$\operatorname{arsinh} x$	$\frac{1}{\sqrt{x^2 + 1}}$	Yes
$\operatorname{arcosh} x$	$\frac{1}{\sqrt{x^2 - 1}} \quad x > 1$	Yes
$\operatorname{artanh} x$	$\frac{1}{1 - x^2} \quad x < 1$	Yes

As with any non-trivial integration, always scan straight away for a possible “Chain Rule Backwards” before reaching for the much bigger weapons in the armoury; Integration by Substitution and Integration by Parts.

8.2 Example

Find $\int \frac{2 + 5x}{\sqrt{x^2 + 1}} dx$

Teaching Video : <http://www.NumberWonder.co.uk/v9102/8.mp4>



[6 marks]

[†] Hyperbolic Functions, Lesson 6

8.3 Exercise

*Any solution based entirely on graphical
or numerical methods is not acceptable*
Marks Available : 50

Question 1

Integrate the following with respect to x ,

(i) $\sinh(5x)$ (ii) $\frac{1}{\cosh^2(4x)}$

[2, 3 marks]

Question 2

Marvin is thinking of using integration by parts twice to find the following integral,

$$\int x^2 \cosh(x^3 + 4) dx$$

but Giles points out that it can far more easily be done as a “chain rule backwards”.
Following Giles' advice, carry out the integration.

[2 marks]

Question 3

By setting up a “chain rule backwards” find,

(i) $\int \sinh(5x) \cosh^6(5x) dx$

(ii) $\int \frac{\sinh(5x)}{\cosh^6(5x)} dx$

[4 marks]

Question 4

Frederic wishes to integrate the $\tanh x$ function and has begun by setting up a “chain rule backwards”.

(i) Complete Fredric's solution,

$$\begin{aligned}\int \tanh x \, dx &= \int \frac{\sinh x}{\cosh x} \, dx \\ &= \int \sinh x (\cosh x)^{-1} \, dx\end{aligned}$$

[2 marks]

(ii) Justify why no modulus signs are needed in the final result.

[1 mark]

Question 5

(i) Use the approach outlined in question 4 to find $\int \coth x \, dx$

[2 marks]

(ii) Are modulus signs are needed in the final result ?

[1 mark]

Question 6

Find the exact value of $\int_0^{\sqrt{2}} \frac{6}{\sqrt{1 + 4x^2}} \, dx$


[5 marks]

Question 7

The graph is of the function $f(x) = \frac{1+x}{\sqrt{x^2+1}}$ (in red) along with the vertical line (in gold) with equation $x = \sqrt{3}$. Show that the exact value of the area bounded by this vertical line, $f(x)$, and the x -axis is,

$$\ln((\sqrt{3} + 2)(\sqrt{2} + 1)) + 2 - \sqrt{2}$$

The techniques of example 8.2 will be useful !



[8 marks]

Question 8

By replacing the $\cosh x$ function with exponentials, find $\int_0^{\ln 3} e^x \cosh(2x) \, dx$

[4 marks]

Question 9


FM A-Level Examination Question from June 2014, Paper FP3, Q3(b) (Edexcel)

Using calculus, find the exact value of $\int_0^1 e^{2x} \sinh x \, dx$

[4 marks]

Question 10

FM A-Level Examination Question from June 2013, Paper FP3, Q7 (Edexcel)



The curves shown are $y = 6 \cosh x$ (red) and $y = 9 - 2 \sinh x$ (gold).

- (a) Using the definitions of $\sinh x$ and $\cosh x$ in terms of e^x , find exact values for the x -coordinate of the two points where the curves intersect.

[6 marks]

The finite region between the two curves is shown shaded.

- (b) Using calculus, find the area of the shaded region, giving your answer in the form $a \ln b + c$, where a , b and c are integers.

[6 marks]

This document is a part of a **Mathematics Community Outreach Project** initiated by Shrewsbury School

It may be freely duplicated and distributed, unaltered, for non-profit educational use

In October 2020, Shrewsbury School was voted “**Independent School of the Year 2020**”

© 2025 Number Wonder

Teachers may obtain detailed worked solutions to the exercises by email from MHHShrewsbury@Gmail.com