# Push The Pace #3

You have thirty-five minutes to answer seven examination questions

Marks Available: 40 (+ 10 bonus)


Further A-Level Pure Mathematics Push The Pace Revision Papers

#### **Question 1**

Further A-Level Examination Question from June 2019, Paper 1, Q4 (OCR)

The graph shows the region bounded by the curve  $y = sec(\frac{x}{2})$ , the x-axis, the

y-axis and the line  $x = \frac{\pi}{2}$ 



This region is rotated through  $2\pi$  radians about the x-axis.

Find, in exact form, the volume of the solid of revolution generated.

Further AS-Level Examination Question from June 2016, Paper FP1. Q1 (CEA)

Let 
$$\mathbf{A} = \begin{pmatrix} 5 & 4 \\ -3 & -2 \end{pmatrix}$$
 and  $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ 

(i) Verify that 
$$A^2 = 3A - 2I$$

[4 marks]

(ii) Hence, or otherwise, express the matrix  $\mathbf{A}^{-1}$  in the form  $\alpha \mathbf{A} + \beta \mathbf{I}$ , where  $\alpha$  and  $\beta$  are real numbers.

Further A-Level Examination Question from June 2019, Paper 1, Q8 (OCR)

The roots of the equation  $x^3 - x^2 + kx - 2 = 0$  are  $\alpha$ ,  $\frac{1}{\alpha}$  and  $\beta$ 

(a) Evaluate, in exact form, the roots of the equation.

[ 6 marks ]

(**b**) Find k

Further AS-Level Examination Question from June 2021, Paper 1, Q8 (AQA) Stephen is correctly told that (1 + i) and -1 are two roots of the polynomial equation  $z^3 - 2iz^2 + pz + q = 0$  where p and q are complex numbers.

(a) Stephen states that (1 - i) must also be a root of the equation because roots of polynomial equations occur in conjugate pairs. Explain why Stephen's reasoning is wrong.

[ 1 mark ]

(**b**) Find p and q

[ 5 marks ]

#### **Question 5**

Further A-Level Examination Question from June 2019, Paper 1, Q4 (AQA) Solve the equation  $2z - 5iz^* = 12$ 

$\sim$	4 •	-
U	uestion	O

Further A-Level Examination Question from June 2015, Paper FP3, Q1 (WJEC)

(a) Express  $5 \cosh \theta + 3 \sinh \theta$  in the form  $r \cosh (\theta + \alpha)$ , r > 0, where the values of r and  $\alpha$  are to be found.

[ 4 marks ]

(**b**) Hence solve the equation  $5 \cosh \theta + 3 \sinh \theta = 10$ 

Further A-Level Examination Question from June 2017, Paper FP3, Q6 (WJEC)

The integral  $I_n$  is given, for  $n \ge 0$ , by  $I_n = \int_0^{\frac{\pi}{4}} tan^n x \, dx$ 

(a) Show that, for 
$$n \ge 2$$
,  $I_n = \frac{1}{n-1} - I_{n-2}$ 

Hint : set up a chain rule backwards keeping in mind that the derivative of tan x is  $sec^2 x$ 

(**b**) Hence determine the value of the integral,  $\int_0^{\frac{\pi}{4}} (3 + tan^2 x)^2 dx$  leaving your answer in terms of  $\pi$